in technological processes, environmental studies, engineering systems, and many others, and the results obtained help to better understand and control heat dissipation.

References:

1. A.A.Samarskii and V.A. Galaktionov, S.P.Kurdyumov, A.P.Mikhailov, Blowe-up in parabolic equations, (Bulletin (new series) of the american mathematical society, 33(4), 1996), pp. 483–486.
2. J.M.Toshtemirov. Effects of a multicomponent heat source on ambient density in multidimensional fields, " ILM SARCHASHMALARI", 2-son (fevral 2025), pp. 33–37.
3. Adrien Drouillet, Romain Le Tellier, Raphael Loubere, Mathieu Peybernes, Louis Viot. Multidimensional simulation of phase change by a D-2D model coupling via Stefan condition. Communications on Applied Mathematics and Computation, 2021, 5 (2), pp. 883–884. ff10.1007/s42967- 021-00157-y. hal-03358560f.
4. Aripov M.M., Djabbarov O.R., Sh.Sadullaeva. Mathematic modeling of processes describing by double nonlinear parabolic equation with convective transfer and damping // AIP Conference Proceedings, 2021, 2365, 060008.
5. A.T.Khaidarov and J.M.Toshtemirov. Modeling of heat propagation processes in multidimensional domains. Modern Probl. of Appl. Math. and Inform. Techn. Al-Khwarizmi, Tashkent, 2024, pp. 60–61.
6. A.T.Khaidarov and J.M.Toshtemirov. Heat source density in non-linear heat dissipation processes. Proceedings of Scientific Conference on Multidisciplinary Studies, Russia, 2023, pp. 72–80.
7. A.Mamatov and J.Toshtemirov. Visualization of the problem of multidimensional heat transfer through digital technologies. Pedagogical reforms and their solutions, Tashkent, 2024, pp. 44–47.
8. A.A.Samarskiy. Teoriya raznostnyx sxem. Nauka. M.: 1989 g. – 616 s.
9. Aripov M.M., Djabbarov O.R., Sh.Sadullaeva. Mathematic modeling of processes describing by double nonlinear parabolic equation with convective transfer and damping. AIP Conference Proceedings, 2021, 2365, 060008.
10. U.U.Begulov, Kh.Abdugappor va J.M.Toshtemirov. Cauchy problem for a parabolic equation describing the heat propagation process in a non-divergent form under the influence of an exponentially varying density. "Matematik fizikaning zamonaviy usullari" ilmiy konferensiyasi va ularning qo'llanilishi, 22.04, 2025

Akhmedov Kurbanbay Yuldashboyevich (Teacher of the Department of Mathematical Analysis at Urgench State University named after Abu Rayhan Beruni; email.qurbanbayaxmedov@urdu.uz},
Bekchanova Shakhnoza Khusanboy qizi (2nd-year master's student in Mathematics, Urgench State University named after Abu Rayhan Beruni; Email: shakhnozabekchanovaa@gmail.com)

ON COMPLEX HYPERPLANES IN $\mathbb{C}^n$ SPACE AND THEIR PROJECTIONS IN REAL SPACE

Annotation. *This article studies the projections of complex hyperplanes in the real space of $\mathbb{C}^n$. In particular, it proves a theorem stating that for any $m \le n$ dimensional real plane taken from the space $\mathbb{R}^n \subset \mathbb{C}^n$, there exists a unique m-dimensional complex plane that contains it.*

Key words*: real space, complex space, real plane, complex hyperplane.*

Axmedov Qurbonbay Yuldashboyevich (Abu Rayhon Beuniy nomidagi Urganch davlat universiteti katta o'qituvchisi; email.qurbonbayaxmedov@urdu.uz),
Bekchanova Shaxnoza Xusanboy qizi (Abu Rayhon Beruniy nomidagi Urganch davlat universiteti Matematika mutaxassisligi 2-kurs magistranti; e-mail.shakhnozabekchanovaa@gmail.com)

$\mathbb{C}^n$ FAZODAGI KOMPLEKS GIPERTEKISLIKLAR VA ULARNING HAQIQIY FAZODAGI PROYEKSIYALARI HAQIDA

Annotatsiya. *Ushbu maqolada $\mathbb{C}^n$ fazodagi kompleks gipertekisliklarning haqiqiy fazodagi proyeksiyalari o'rganilgan bo'lib, xususan, $\mathbb{R}^n \subset \mathbb{C}^n$ fazodan olingan ixtiyoriy $m \le n$ o'lchovli haqiqiy tekislikni o'z ichiga oluvchi yagona m o'lchovli kompleks tekislik mavjudligi haqidagi teorema isbotlangan.*

Kalit so'zlar: *haqiqiy fazo, kompleks fazo, haqiqiy tekislik, kompleks gipertekislik.*

Ахмедов Курбанбай Юлдашбоевич (старший преподаватель Ургенчского государственного университета имени Абу Райхана Беруни; email.qurbanbayaxmedov@urdu.uz),
Бекчанова Шахноза Хусанбойевна (2-й курс магистратуры по математике Ургенчского государственного университета имени Абу Райхана Беруни; email.shakhnozabekchanovaa@gmail.com)

О КОМПЛЕКСНЫХ ГИПЕРПЛОСКОСТЯХ В $\mathbb{C}^n$ ПРОСТРАНСТВЕ И ИХ ПРОЕКЦИЯХ В РЕАЛЬНОМ ПРОСТРАНСТВЕ

Аннотация. *В данной статье исследуются проекции комплексных гиперплоскостей в вещественном пространстве $\mathbb{C}^n$. В частности, доказывается теорема, утверждающая, что для любой $m \le n$ мерной вещественной плоскости, взятой из пространства $\mathbb{R}^n \subset \mathbb{C}^n$, существует единственная m-мерная комплексная плоскость, содержащая её.*

Ключевые слова: *вещественное пространство, комплексное пространство, вещественная плоскость, комплексная гиперплоскость.*

Introduction. It is known that functions possessing subharmonic properties on the sections of m-dimensional planes in an $n-$ dimensional real space are among the important objects of modern geometry in $\mathbb{R}^n$. In this direction, significant results have been obtained in the works of R.Harvey and B.Lawson [4-9], B.Drnovsek, B.D. Forstnerich [3], as well as other mathematicians. These papers consider a class of functions that are subharmonic on $p-$dimensional real planes in the Euclidean space $\mathbb{R}^n$. This class of functions is called $p-$plurisubharmonic functions, which were later successfully applied in $p-$convex geometry to describe the $p-$convex hulls of geometric objects.

In recent years, studying such a class of functions in $\mathbb{C}^n$, in the complex space on m-dimensional complex hyperplanes, through their subharmonic properties, has become one of the leading research directions. This, in turn, raises the problem of studying the relationship between complex hyperplanes and real planes. The results obtained in this paper, namely the study of the projections of complex hyperplanes in real space, play a significant role in solving such problems.

Research Methodology. m-dimensional hyperplanes in $\mathbb{R}^n$. (see. [2]). The $m-$dimensional hyperplanes are represented by the following equations:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \ldots + a_{1n}x_n = \beta_1 \\ a_{21}x_1 + a_{22}x_2 + \ldots + a_{2n}x_n = \beta_2 \\ \ldots \\ a_{n-m,1}x_1 + a_{n-m,2}x_2 + \ldots + a_{n-m,n}x_n = \beta_{n-m}. \end{cases} \tag{1}$$

Here $x_1, x_2, \ldots, x_n$ are the coordinates of a point in $\mathbb{R}^n$, $a_{ij} \in \mathbb{R}^n$ are the coefficients representing the orientation of each hyperplane, $\beta_{ij} \in \mathbb{R}^n$ are constants determining the distance of each hyperplane from the origin. To define an $m-$dimensional hyperplane in $\mathbb{R}^n$, the system must consist of exactly $n-m$ **linearly independent** equations.

m-dimensional real and complex hyperplanes in $\mathbb{C}^n$.(see. [1]) Now suppose we are given $k \le 2n$ vectors $a^\mu \in \mathbb{C}^n$ that are linearly independent over $\mathbb{R}$; the set of points $z \in \mathbb{C}^n$ that satisfies the following system of k real equations

Let us assume that we have $k \le 2n$ vectors $a^\mu \in \mathbb{C}^n$ that are linearly independent over $\mathbb{R}$. We consider the set of points $z \in \mathbb{C}^n$ that satisfy the following system of k real equations.

$$\mathrm{Re}(z, a^\mu) = \beta_\mu \quad \mu = 1, \ldots, k, \tag{2}$$

is known as a real plane of codimension k or a plane of dimension $m = 2n - k$ (for $k = 2n$, this is a point). Since k represents the number of independent constraints imposed on the system, the resulting dimension m effectively reflects the degrees of freedom remaining for the points on the plane.

If the vectors a^μ are instead linearly independent over $\mathbb{C}$ (thus requiring $k \le n$, as k cannot exceed the dimension of the complex space), then the set described by the system of complex equations

$$(z, a^\mu) = b_\mu, \ \mu = 1, \ldots, k, \tag{3}$$

is called a complex plane of codimension k or dimension $m = n - k$. It follows that such a plane is determined by both the complex constraints imposed by a^μ and the specific values b_μ. Since this system can be rewritten as $2k$ real linear equations $\mathrm{Re}(z, a^\mu) = \mathrm{Re}\, b_\mu$, $\mathrm{Re}(z, ia^\mu) = \mathrm{Im}\, b_\mu$ and the vectors a^μ and ia^μ ($\mu = 1, \ldots, k$) are linearly independent over $\mathbb{R}$, then this is a plane of real codimension $2k$.

Building on the foundational concepts of R^n and $\mathbb{C}^n$ spaces, we now present the main result of this study. The following theorem shows the relationship between m-dimensional hyperplanes of $\mathbb{R}^n$ and $\mathbb{C}^n$ where $\mathbb{R}^n_x \subset \mathbb{C}^n_z = \mathbb{R}^n_x + i\mathbb{R}^n_y \left(z = x + iy\right)$

Analysis and results. Theorem. In the space $\mathbb{C}^n$, a real m-dimensional hyperplane $\prod$ in $\mathbb{R}^n_x \subset \mathbb{C}^n$ is contained within an unique m-dimensional complex hyperplane.

Proof. Let $\prod$ be a real m-dimensional plane ($\dim \prod = m$) in $\mathbb{R}^n_x$. The given m-dimensional real plane is defined by the following system of equations:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \ldots + a_{1n}x_n = \beta_1 \\ a_{21}x_1 + a_{22}x_2 + \ldots + a_{2n}x_n = \beta_2 \\ \ldots \\ a_{n-m,1}x_1 + a_{n-m,2}x_2 + \ldots + a_{n-m,n}x_n = \beta_{n-m}. \end{cases} \tag{4}$$

Here a_{jk} and β_j are real numbers ($j = \overline{1, n-m}$).

We are given the complex space which can be written as $\mathbb{C}^n = \mathbb{R}^n_x + i\mathbb{R}^n_y$, where m-dimensional complex plane $\dim \prod^{\mathbb{C}} = m$ is given. Let this complex plane defined by the following system of equations:

$$\begin{cases} c_{11}z_1 + c_{12}z_2 + \ldots + c_{1n}z_n = \gamma_1 \\ c_{21}z_1 + c_{22}z_2 + \ldots + c_{2n}z_n = \gamma_2 \\ \ldots \\ c_{n-m,1}z_1 + c_{n-m,2}z_2 + \ldots + c_{n-m,n}z_n = \gamma_{n-m}. \end{cases} \tag{5}$$

Here c_{jk} and γ_j are complex numbers ($j = \overline{1, n-m}$).

Systems (4) and (5) can be written in the following form:

$$a_{j1}x_1 + a_{j2}x_2 + \ldots + a_{jn}x_n = \beta_j,$$

$$c_{j1}z_1 + c_{j2}z_2 + \ldots + c_{jn}z_n = \gamma_j.$$

where $j = \overline{1, n-m}$. $\operatorname{Re}(c_{j1}z_1 + c_{j2}z_2 + \ldots + c_{jn}z_n) = \operatorname{Re}\gamma_j$ is equivalent to $a_{j1}x_1 + a_{j2}x_2 + \ldots + a_{jn}x_n = \beta_j$ according to the construction of the space $\mathbb{C}^n$.

$$\operatorname{Re}\left(\sum_{k=1}^{n} c_{jk}z_k\right) = \operatorname{Re}\left(\sum_{k=1}^{n} c_{jk}(x_k + iy_k)\right) = \sum_{k=1}^{n}\left(\operatorname{Re}c_{jk}x_k - \operatorname{Im}c_{jk}y_k\right) = \operatorname{Re}\gamma_j$$

$$\begin{cases} \sum_{k=1}^{n}\left(\operatorname{Re}c_{jk}x_k - \operatorname{Im}c_{jk}y_k\right) = \operatorname{Re}\gamma_j \\ a_{j1}x_1 + a_{j2}x_2 + \ldots + a_{jn}x_n = \beta_j \end{cases}$$

to ensure that this equation is consistent, we need to equate the corresponding coefficients. For this, $\operatorname{Re}c_{jk} = a_{jk}$, $\operatorname{Im}c_{jk} = 0$ ($k = \overline{1,n}$ $j = \overline{1, n-m}$ conditions must hold. Thus, $c_{jk} = a_{jk}$ is valid.

From this, we write the following equation:

$$a_{j1}z_1 + a_{j2}z_2 + \ldots + a_{jn}z_n = \gamma_j.$$

Here, $z_n = x_n + iy_n$. Taking this into account:

$$a_{j1}(x_1 + iy_1) + a_{j2}(x_2 + iy_2) + \ldots + a_{jn}(x_n + iy_n) = \gamma_j$$

$$a_{j1}x_1 + a_{j2}x_2 + \ldots + a_{jn}x_n + i\sum_{k=1}^{n} a_{jk}y_k = \gamma_j$$

$$\beta_j + i\sum_{k=1}^{n} a_{jk}y_k = \gamma_j. \tag{6}$$

Consequently, (4) aligns with (5) as: In (6), when considering the real part of the equation $i\sum_{k=1}^{n} a_{jk} y_k = 0$, it follows that $\beta_j = \gamma_j$, ensuring the consistency of the equation. This proves that the coefficients of (4) consist entirely of real numbers.

Examples of Complex Hyperplanes Containing Real Planes in $\mathbb{C}^3$. The theorem states that any real m-dimensional hyperplane in $\mathbb{R}_x^n \subset \mathbb{C}^n$ is uniquely contained within an m-dimensional complex hyperplane. In this section, we provide explicit examples following the structure of the theorem and its proof.

A real plane in $\mathbb{R}^3$ is given by the equation: $a_1 x_1 + a_2 x_2 + a_3 x_3 = \beta$ where a_j and β are real numbers. For instance, consider: $x_1 + 2x_2 - x_3 = 4$. This equation defines a two-dimensional real hyperplane in $\mathbb{R}^3$.

We extend the equation by allowing x_j to take complex values, defining a complex hyperplane in $\mathbb{C}^3$: $z_1 + 2z_2 - z_3 = 4$ where $z_j = x_j + iy_j$ are complex variables.

Expanding in terms of real and imaginary components:

$$(x_1 + iy_1) + 2(x_2 + iy_2) - (x_3 + iy_3) = 4.$$

Separating real and imaginary parts gives the system:

–**Real part:** $x_1 + 2x_2 - x_3 = 4$;

–**Imaginary part:** $y_1 + 2y_2 - y_3 = 0$.

Since the imaginary part is zero, this ensures the given real plane is uniquely contained in the complex hyperplane.

Conclusion. This paper examines a special case of the projections of complex hyperplanes in $\mathbb{C}^n$ space onto real space. The obtained result plays an important role in studying the relationships between complex hyperplanes and real planes.

References:

(1). B.V.Shabat. Introduction to complex Analysis Part II Functions of Several variables. American Mathematical Society, (1992), p. 1–12.

(2). Rudin W. Principles of Mathematical Analysis. R.R.Donnelley & Sons. 1976, p. 16–19.

(3). Drnovsek, B.D., Forstnerich F. Minimal hulls of compact sets in $\mathbb{R}^3$. Trans. Am. Math. Soc. 368 (10), p. 7477–7506, October 2016.

(4). Harvey F.R. and Lawson H.B.Jr., An introduction to potential theory in calibrated geometry. Amer. J. Math., V. 131, № 4, (2009), p. 893–944.

(5). Harvey F.R. and Lawson H.B.Jr., Duality of positive currents and plurisubharmonic functions in calibrated geometry. Amer. J. Math.V. 131, № 5, (2009), p. 1211–1240.

(6). Harvey F.R. and Lawson H.B.Jr., Geometric plurisubharmonicity and convexity – an introduction. Advances in Mathematics. V. 230, (2012), p. 2428–2456.

(7). Harvey F.R. and Lawson H.B.Jr., Plurisubharmonicity in a general geometric context. Geometry and Analysis I, (2010), p. 363–401.

(8). Harvey F.R. and Lawson H.B.Jr., Calibrated geometries. Acta Mathematica 148, 1982, p. 47–157.

(9). Harvey F.R. and Lawson H.B.Jr., $p-$convexity, $p-$plurisubharmonicity and the Levi problem// Indiana Univ. Math. J. 62, № 1 (2013), p. 149–169.

(10). Joyce D. Riemannian holonomy groups and calibrated geometry. Oxford University Press, USA. 2007, 318 p.